

ALPHA-AMYLASE (Bacillus licheniformis) (Lot 120901a)

E-BLAAM-100ML

08/15

(EC 3.2.1.1) 4-alpha-D-glucan glucanohydrolase CAZy: GH Family 13

PROPERTIES

I. ELECTROPHORETIC PURITY:

- Single major band on isoelectric focusing (pl = 7.4)
- Single major band on SDS-gel electrophoresis (MW = 58,000)

2. SPECIFIC ACTIVITY AND LEVEL OF OTHER ACTIVITIES:

| Substrate | Specific Activity (U/mg Protein) |
|---|-------------------------------------|
| α -Amylase (Ceralpha Reagent at pH 6.0) | 54.0 |
| Amyloglucosidase (p-Nitrophenyl β -maltoside) | undetectable |
| Cellulase (CM-Cellulose 4M) | undetectable |
| β -Mannanase (carob galactomannan) | undetectable |

One Unit of α -amylase is the amount of enzyme required to release one μ mole of *p*-nitrophenol from blocked *p*-nitrophenyl-maltoheptaoside per minute (in the presence of excess α -glucosidase) at pH 6.0 and 40°C.

3. PHYSICOCHEMICAL PROPERTIES:

| pH Optima: | 6.0-6.5 |
|------------------------|---------|
| pH Stability: | 4.5-8.0 |
| Temperature Optima: | 75°C |
| Temperature Stability: | < 80°C |

4. **STORAGE CONDITIONS:**

The enzyme is supplied in vials of 100 mL as a stabilised solution and should be stored at $4^{\circ}C$.

The enzyme is supplied at a concentration of 3000 U/mL on Ceralpha Reagent at pH 6.0 and 40°C (i.e. approximately 10,000 U/mL on soluble starch under the same assay conditions).

This enzyme is recommended for use in **Total Dietary Fibre** analytical procedures and the **Megazyme Total Starch** test method. The preparation is effectively devoid of cellulase and is free of catalase.