



[www.megazyme.com](http://www.megazyme.com)

---

## **FORMIC ACID (FORMATE)**

### **ASSAY PROCEDURE**

K-FORM 10/20

(\*25 Manual Assays per Kit) or  
(220 Auto-Analyser Assays per Kit) or  
(250 Microplate Assays per Kit)

*\*The number of tests per kit can be doubled if all volumes are halved*



## INTRODUCTION:

Formic acid is a commonly occurring intermediate in cellular metabolism and is excreted in urine as a normal physiological product. Formic acid is produced as a degradation product of certain amino acids, e.g. methionine, serine and glycine, as well as of several organic substances such as methanol and acetone.

Formic acid is a weak, colourless, volatile organic acid ( $\text{H}_2\text{CO}_2$ ). It occurs naturally in ants and in the fruit of the soap tree, and is also formed as a by-product in the atmospheric oxidation of turpentine. The principal commercial product is sodium formate, which is prepared by the reaction of carbon monoxide and sodium hydroxide under pressure and heat. Formic acid is used in leather manufacture to control pH, as well as in acid dyeing.

The principal use of formic acid is as a preservative (E236). It is used as an antibacterial agent in livestock feed. When sprayed on fresh hay or other silage, it arrests certain decay processes and causes the feed to retain its nutritive value longer, and so it is widely used to preserve winter feed for cattle. In the poultry industry, it is sometimes added to feed to kill salmonella bacteria. As a preservative in fruit juices and pickles it has an antiseptic effect against yeasts. Some beekeepers use formic acid as a miticide against the Varroa mite. Some formate esters are artificial flavourings or perfumes.

## PRINCIPLE:

In the presence of  $\text{NAD}^+$ , formic acid is oxidised to carbon dioxide ( $\text{CO}_2$ ) by the enzyme formate dehydrogenase (FDH) (I) with the concurrent formation of NADH.



The amount of NADH formed is stoichiometric with the amount of formic acid. It is the NADH which is measured by the increase in absorbance at 340 nm.

## SPECIFICITY, SENSITIVITY, LINEARITY AND PRECISION:

The method is specific for formic acid. Acetic acid, propionic acid, ascorbic acid and oxalic acid do not react.

The smallest differentiating absorbance for the assay is 0.005 absorbance units. This corresponds to 0.0466 mg/L of sample solution at the maximum sample volume of 2.00 mL (or 0.932 mg/L with a sample volume of 0.1 mL). The detection limit is 0.0932 mg/L, which is derived from an absorbance difference of 0.01 with the maximum sample volume of 2.00 mL.

Formic acid is split into carbon dioxide and water during storage and thus in the analysis of commercial formic acid recoveries of < 100% can be expected. The volatility of the solution must also be taken into consideration.

The assay is linear over the range of 0.4 to 20 µg of formic acid per assay. In duplicate determinations using one sample solution, an absorbance difference of 0.005 to 0.010 may occur. With a sample volume of 2.00 mL, this corresponds to a formic acid concentration of approx. 0.0466 to 0.0932 mg/L. If the sample is diluted during sample preparation, the result is multiplied by the dilution factor, F. If, in sample preparation, the sample is weighed, e.g. 10 g/L, a difference of 0.02 to 0.05 g/100g can be expected.

### **INTERFERENCE:**

If the conversion of formic acid has been completed within the specified time (approx. 12 min), it can generally be concluded that no interference has occurred. However, this can be further checked by adding formic acid (approx. 10 µg in 0.1 mL) to the cuvette on completion of the reaction. A significant increase in the absorbance should be observed.

Interfering substances in the sample being analysed can be identified by including an internal standard. Quantitative recovery of this standard would be expected. Losses in sample handling and extraction are identified by performing recovery experiments, i.e. by adding formic acid to the sample in the initial extraction steps.

Formate dehydrogenase is inhibited by sodium azide, so this should be excluded from all assay components and samples.

### **SAFETY:**

The general safety measures that apply to all chemical substances should be adhered to.

For more information regarding the safe usage and handling of this product please refer to the associated SDS that is available from the Megazyme website.

## KITS:

Kits suitable for performing 25 assays in manual format (or 220 assays in auto-analyser format or 250 assays in microplate format) are available from Megazyme. The kits contain the full assay method plus:

**Bottle 1: (x2)** Buffer (freeze-dried powder, pH 7.6).  
Stable for > 2 years at 4°C.

**Bottle 2:** NAD<sup>+</sup>.  
Stable for > 2 years below -10°C.

**Bottle 3:** Formate dehydrogenase (1.4 mL).  
Stable for > 2 years at 4°C.

**Bottle 4:** Sodium formate (~ 2 g).  
Stable for > 2 years at 4°C.

## PREPARATION OF REAGENT SOLUTIONS/SUSPENSIONS:

1. Dissolve the contents of one of bottle 1 in 4 mL of distilled water. Store below -10°C between use. Stable for > 2 years below -10°C. Do not dissolve the contents of the second bottle until required.
2. Dissolve the contents of bottle 2 in 5.2 mL of distilled water. Divide into appropriately sized aliquots and store in polypropylene tubes below -10°C between use and keep cool during use if possible. Once dissolved, the reagent is stable for > 2 years below -10°C.
3. Use the contents of bottle 3 as supplied. Before opening for the first time, shake the bottle to remove any enzyme that may have settled on the rubber stopper. Subsequently, store the bottle in an upright position. **Swirl the bottle to mix contents before use.**  
Stable for > 2 years at 4°C.
4. Accurately weigh 148 mg of sodium formate (MW = 68.08) to the nearest 0.1 mg into a 1 L volumetric flask. Fill to the mark with distilled water and mix thoroughly. This corresponds to 0.1 g/L formic acid. Divide into appropriately sized aliquots and store below -10°C.  
Stable for 1 year below -10°C.

**NOTE:** The sodium formate standard solution is only assayed where there is some doubt about the accuracy of the spectrophotometer being used or where it is suspected that inhibition is being caused by substances in the sample. The concentration of formic acid is determined directly from the extinction coefficient of NADH (see page 6).

## EQUIPMENT (RECOMMENDED):

1. Glass test tubes (round bottomed; 16 x 100 mm).
2. Disposable plastic cuvettes (1 cm light path, 3.0 mL).
3. Micro-pipettors, e.g. Gilson Pipetman<sup>®</sup> (20  $\mu$ L, 100  $\mu$ L and 200  $\mu$ L).
4. Positive displacement pipettor, e.g. Eppendorf Multipette<sup>®</sup>
  - with 5.0 mL Combitip<sup>®</sup> (to dispense 0.2 mL aliquots of NAD<sup>+</sup> solution).
  - with 25 mL Combitip<sup>®</sup> (to dispense 2.0 mL aliquots of distilled water).
5. Analytical balance.
6. Spectrophotometer set at 340 nm.
7. Vortex mixer (e.g. IKA<sup>®</sup> Yellowline Test Tube Shaker TTS2).
8. Stop clock.
9. Whatman No. 1 (9 cm) filter papers.

## A. MANUAL ASSAY PROCEDURE:

**Wavelength:** 340 nm  
**Cuvette:** 1 cm light path (glass or plastic)  
**Temperature:** ~ 25°C  
**Final volume:** 2.55 mL  
**Sample solution:** 0.4-20 µg of formic acid per cuvette  
(in 0.10-2.00 mL sample volume)

**Read against air** (without a cuvette in the light path) or against water

Pipette into cuvettes	Blank	Sample
distilled water (at ~ 25°C)	2.10 mL	2.00 mL
sample*	-	0.10 mL
solution 1 (buffer)	0.20 mL	0.20 mL
solution 2 (NAD <sup>+</sup> )	0.20 mL	0.20 mL
Mix**, read the absorbances of the solutions ( $A_1$ ) after approx. 5 min and start the reactions by addition of:		
suspension 3 (FDH)	0.05 mL	0.05 mL
Mix** and read the absorbances of the solutions ( $A_2$ ) at the end of the reaction (approx. 12 min).		

\* rinse the dispensing pipette tip with the solution before dispensing the sample.

\*\* for example with a plastic spatula or by gentle inversion after sealing the cuvette with a cuvette cap or Parafilm®.

## CALCULATION:

Determine the absorbance difference ( $A_2 - A_1$ ) for both blank and sample. Subtract the absorbance difference of the blank from the absorbance difference of the sample, thereby obtaining

$\Delta A_{\text{formic acid}}$ . The value of  $\Delta A_{\text{formic acid}}$  should as a rule be at least 0.100 absorbance units to achieve sufficiently accurate results.

The concentration of formic acid can be calculated as follows:

$$c = \frac{V \times MW}{\varepsilon \times d \times v} \times \Delta A_{\text{formic acid}} \quad [\text{g/L}]$$

### where:

V = final volume [mL]

MW = molecular weight of formic acid [g/mol]

$\varepsilon$  = extinction coefficient of NADH at 340 nm  
= 6300 [ $\text{l} \times \text{mol}^{-1} \times \text{cm}^{-1}$ ]

d = light path [cm]

v = sample volume [mL]

### It follows for formic acid:

$$c = \frac{2.55 \times 46.03}{6300 \times 1.0 \times 0.1} \times \Delta A_{\text{formic acid}} \quad [\text{g/L}]$$
$$= 0.1863 \times \Delta A_{\text{formic acid}} \quad [\text{g/L}]$$

If the sample has been diluted during preparation, the result must be multiplied by the dilution factor, F.

When analysing solid and semi-solid samples which are weighed out for sample preparation, the content (g/100 g) is calculated from the amount weighed as follows:

### Content of formic acid

$$= \frac{C_{\text{formic acid}} [\text{g/L sample solution}]}{\text{weight}_{\text{sample}} [\text{g/L sample solution}]} \times 100 \quad [\text{g/100 g}]$$

**NOTE:** These calculations can be simplified by using the Megazyme **Mega-Calc™**, downloadable from where the product appears on the Megazyme website ([www.megazyme.com](http://www.megazyme.com)).

## B. AUTO-ANALYSER ASSAY PROCEDURE:

### NOTES:

1. The Auto-Analyser Assay Procedure for formic acid can be performed using either a single point standard or a full calibration curve.
2. For each batch of samples that is applied to the determination of formic acid **either a single point standard or a calibration curve must be performed concurrently using the same batch of reagents.**

Reagent preparation is performed as follows:

### Preparation of R1:

Component	Volume
distilled water	22.2 mL
solution 1 (buffer)	2.5 mL
solution 2 (NAD <sup>+</sup> )	2.5 mL (after adding 5.2 mL of H <sub>2</sub> O to bottle 2)
Total volume	27.2 mL

### Preparation of R2:

Component	Volume
distilled water	2.30 mL
suspension 3 (FDH)	0.51 mL
Total volume	2.81 mL

### EXAMPLE METHOD:

**R1:** 0.200 mL

Sample: ~ 0.01 mL

**R2:** 0.025 mL

**Reaction time:** ~ 12 min at 37°C

**Wavelength:** 340 nm

**Prepared reagent stability:** > 2 days when refrigerated

**Calculation:** endpoint

**Reaction direction:** increase

**Linearity:** up to 0.183 g/L of formic acid using 0.01 mL sample volume

## C. MICROPLATE ASSAY PROCEDURE:

### NOTES:

1. The Microplate Assay Procedure for formic acid can be performed using either a single point standard or a full calibration curve.
2. For each batch of samples that is applied to the determination of formic acid **either a single point standard or a calibration curve must be performed concurrently using the same batch of reagents.**

**Wavelength:** 340 nm  
**Microplate:** 96-well (e.g. clear flat-bottomed, glass or plastic)  
**Temperature:** ~ 25°C  
**Final volume:** 0.255 mL  
**Linearity:** 0.04-2.0 µg of formic acid per well  
(in 0.01-0.20 mL sample volume)

Pipette into wells	Blank	Sample	Standard
distilled water	0.210 mL	0.200 mL	0.200 mL
sample solution	-	0.010 mL	-
standard solution	-	-	0.010 mL
solution 1 (buffer)	0.020 mL	0.020 mL	0.020 mL
solution 2 (NAD <sup>+</sup> )	0.020 mL	0.020 mL	0.020 mL
Mix*, read the absorbances of the solutions (A <sub>1</sub> ) after approx. 5 min and start the reactions by addition of:			
suspension 3 (FDH)	0.005 mL	0.005 mL	0.005 mL
Mix* and read the absorbances of the solutions (A <sub>2</sub> ) at the end of the reaction (approx. 12 min). If the reaction has not stopped after 12 min, continue to read the absorbances at 2 min intervals until the absorbances increase constantly over 2 min**.			

\* for example using microplate shaker, shake function on a microplate reader or repeated aspiration (e.g. using a pipettor set at 50-100 µL volume).

\*\* if this “creep” rate is greater for the sample than for the blank, extrapolate the sample absorbances back to the time of addition of suspension 3.

### CALCULATION (Microplate Assay Procedure):

$$\text{g/L} = \frac{\Delta A_{\text{sample}}}{\Delta A_{\text{standard}}} \times \text{g/L standard} \times F$$

If the sample is diluted during preparation, the result must be multiplied by the dilution factor, F.

## SAMPLE PREPARATION:

### 1. Sample dilution.

The amount of formic acid present in the cuvette (i.e. in the 0.1 mL of sample being analysed) should range between 0.4 and 20  $\mu\text{g}$ . The sample solution must therefore be diluted sufficiently to yield a formic acid concentration between 0.004 and 0.20 g/L.

**Dilution Table**

Estimated concentration of formic acid (g/L)	Dilution with water	Dilution factor (F)
< 0.2	No dilution required	1
0.2-2.0	1 + 9	10
2.0-20	1 + 99	100
> 20	1 + 999	1000

If the value of  $\Delta A_{\text{formic acid}}$  is too low (e.g. < 0.100), weigh out more sample or dilute less strongly. Alternatively, the sample volume to be pipetted into the cuvette can be increased up to 2.00 mL, making sure that the sum of the sample and distilled water components in the reaction is 2.10 mL and using the new sample volume in the equation.

### 2. Sample clarification.

#### a. Solutions:

**Carrez I solution.** Dissolve 3.60 g of potassium hexacyanoferrate (II)  $\{\text{K}_4[\text{Fe}(\text{CN})_6] \cdot 3\text{H}_2\text{O}\}$  (Sigma cat. no. P9387) in 100 mL of distilled water. Store at room temperature.

**Carrez II solution.** Dissolve 7.20 g of zinc sulphate  $(\text{ZnSO}_4 \cdot 7\text{H}_2\text{O})$  (Sigma cat. no. Z4750) in 100 mL of distilled water. Store at room temperature.

**Sodium hydroxide (NaOH, 100 mM).** Dissolve 4 g of NaOH in 1 L of distilled water. Store at room temperature.

#### b. Procedure:

Pipette the liquid sample into a 100 mL volumetric flask which contains approx. 60 mL of distilled water, or weigh sufficient quantity of the sample into a 100 mL volumetric flask and add 60 mL of distilled water. Carefully add 5 mL of Carrez I solution, 5 mL of Carrez II solution and 10 mL of NaOH solution (100 mM). Mix after each addition. Fill the volumetric flask to the mark, mix and filter.

### 3. General considerations.

**(a) Liquid samples:** clear, slightly coloured and approximately neutral, liquid samples can be used directly in the assay.

**(b) Acidic samples:** if  $> 0.1$  mL of an acidic sample is to be used undiluted (such as wine or fruit juice), the pH of the solution should be increased to approx. 8.0 using 2 M NaOH, and the solution incubated at room temperature for 30 min.

**(c) Carbon dioxide:** samples containing significant quantities of carbon dioxide, such as beer, should be degassed by increasing the pH to approx. 8.0 with 2 M NaOH and gentle stirring, or by stirring with a glass rod.

**(d) Coloured samples:** an additional sample blank, i.e. sample with no FDH, may be necessary in the case of coloured samples.

**(e) Strongly coloured samples:** if used undiluted, strongly coloured samples should be treated by the addition of 0.2 g of PVPP/10 mL of sample. Shake the tube vigorously for 5 min and then filter through Whatman No. 1 filter paper.

**(f) Solid samples:** homogenise or crush solid samples in distilled water and filter if necessary.

**(g) Samples containing fat:** extract such samples with hot water at a temperature above the melting point of the fat, e.g. in a 100 mL volumetric flask. Adjust to room temperature and fill the volumetric flask to the mark with distilled water. Store on ice or in a refrigerator for 15-30 min and then filter. Discard the first few mL of filtrate, and use the clear supernatant (which may be slightly opalescent) for assay. Alternatively, clarify with Carrez reagents.

**(h) Samples containing protein:** deproteinise samples containing protein by adding an equal volume of ice-cold 1 M perchloric acid with mixing. Centrifuge at 1,500 g for 10 min and neutralise the supernatant with 1 M KOH. Alternatively, use Carrez reagents.

## **SAMPLE PREPARATION EXAMPLES:**

### **(a) Determination of formic acid in wine.**

For red wine, add 100 mg of charcoal to 5 mL of sample in a polypropylene tube. Mix by inversion for 1 min and then filter the suspension through Whatman GF/A glass fibre filter paper. For white wine, analyse the sample with no prior pre-treatment. *Typically, no dilution is required and a sample volume of 0.1 mL is satisfactory.*

### **(b) Determination of formic acid in fruit juice.**

For strongly coloured juices, add 100 mg of charcoal to 5 mL of sample in a polypropylene tube. Mix by inversion for 1 min and then filter the suspension through Whatman GF/A glass fibre filter paper. Adjust strongly acidic juices to pH 7-8 using 1 M potassium hydroxide. *Typically, no dilution is required and a sample volume of 0.1 mL is satisfactory.*

### **(c) Determination of formic acid in vinegar.**

Add 50 mL of vinegar to a 200 mL beaker and adjust to pH 7-8 with 1 M potassium hydroxide. Quantitatively transfer to a 100 mL volumetric flask and adjust to the mark with distilled water. *Typically, no further dilution is required and a sample volume of 0.2 mL is satisfactory.*

### **(d) Determination of formic acid in pickles.**

Separate liquid from solid by filtration. Store the solution at 4°C for 20 min to obtain separation of fat (if necessary). Filter an aliquot of the aqueous layer, discarding the first few mL. Dilute an aliquot of the filtrate according to the dilution table, if necessary. *Typically, no dilution is required and a sample volume of 0.1 mL is satisfactory.*

### **(e) Determination of formic acid in fruit and vegetable products.**

Homogenise approx. 50 g of fruit or vegetable with 100 mL of water using an electric blender. Stir the mixture for approx. 15 min and quantitatively transfer to a 250 mL volumetric flask and adjust to the mark with distilled water. Mix well. *Typically, no dilution is required and a sample volume of 0.2 mL is satisfactory.*

### **(f) Determination of formic acid in fish and meat products.**

Accurately weigh approx. 5 g of representative homogenised sample into a 50 mL beaker. Add 20 mL perchloric acid solution (1 M) and, using a Polytron® homogeniser or similar, homogenise for 10 min. Adjust the pH to approx. 8 with 2 M potassium hydroxide solution. Quantitatively transfer the mixture to a 100 mL volumetric flask and adjust to the mark with distilled water, ensuring the fatty layer is “above” the mark and that the aqueous layer is “at” the mark. Filter the solution through Whatman No. 1 (9 cm) filter paper. Discard the first few mL and use the clear or slightly turbid solution for the assay. For calculation of the amount of formic acid, take the volume displacement factor of 0.98 into account. *Typically, no dilution is required and a sample volume of 0.5 mL is satisfactory.*

### **(g) Determination of formic acid in bakery goods.**

Accurately weigh 5 g of homogenised or milled sample into a 100 mL volumetric flask. Add approx. 75 mL of distilled water and extract at 20-25°C for 15 min. Fill to the mark with distilled water and filter the solution through Whatman No. 1 (9 cm) filter paper. *Typically, no dilution is required and a sample volume of 0.2 mL is satisfactory.*

### **(h) Determination of formic acid in honey.**


Weigh approx. 10 g of honey into a 100 mL volumetric flask. Fill to the mark with distilled water and mix thoroughly. Use the solution directly in the assay. *Typically, no dilution is required and a sample volume of 0.2 mL is satisfactory.*

### (i) Determination of formic acid in jam.

Accurately weigh approx. 20 g of homogenised sample into a 100 mL beaker. Add approx. 20 mL of hot water ( $\sim 60^{\circ}\text{C}$ ) and mix. Then add 2 mL of Carrez I solution and 2 mL of Carrez II solution. Mix after each addition. Neutralise the sample with 4 mL of 100 mM NaOH. Cool the solution to  $20\text{--}25^{\circ}\text{C}$ , quantitatively transfer to a 100 mL volumetric flask and fill to mark with distilled water. Mix thoroughly and filter through Whatman No. 1 (9 cm) filter paper. Typically, no dilution is required and a sample volume of 0.2 mL is satisfactory.

### (j) Determination of formic acid in protein containing samples.

Add 40 mL of 30 mM trichloroacetic acid to 20 g of protein-containing sample and stir the mixture for 1 min. Neutralise the solution with 1 M potassium hydroxide and quantitatively transfer the mixture to a 100 mL volumetric flask. Fill to mark with distilled water and filter an aliquot through Whatman No. 1 (9 cm) filter paper. Use the clear solution, diluted if necessary, in the assay. Typically, no dilution is required and a sample volume of 0.1 mL is satisfactory.



**Figure 1.** Increase in absorbance at 340 nm on incubation of  $10\ \mu\text{g}$  of formic acid with formate dehydrogenase in the presence of  $\text{NAD}^{+}$ .

### REFERENCE:

Schaller, K. -H. & Triebig, G. (1988). Formate: Determination with Formate Dehydrogenase. "Methods of Enzymatic Analysis" (Bergmeyer, H. U., ed.), 3rd ed., **Vol VI**, pp. 668-672, VCH Publishers (UK) Ltd., Cambridge, UK.

**NOTES:**

[illegible]

**NOTES:**

[illegible]



**Bray Business Park, Bray,  
Co. Wicklow,  
A98 YV29,  
IRELAND.**

**Telephone: (353.1) 286 1220  
Facsimile: (353.1) 286 1264  
Internet: [www.megazyme.com](http://www.megazyme.com)  
E-Mail: [info@megazyme.com](mailto:info@megazyme.com)**

---

---

**WITHOUT GUARANTEE**

The information contained in this booklet is, to the best of our knowledge, true and accurate, but since the conditions of use are beyond our control, no warranty is given or is implied in respect of any recommendation or suggestions which may be made or that any use will not infringe any patents.